

# AU PRODUCT INFORMATION SHEET

Information Provided for Veterinary Professionals Only

## ILIUM MELOXICAM 20

ANTI-INFLAMMATORY INJECTION  
FOR CATTLE, SHEEP, PIGS AND HORSES



## WHAT IS ILIUM MELOXICAM 20 ANTI-INFLAMMATORY INJECTION?

Ilium Meloxicam 20 Anti-Inflammatory Injection is a non-steroidal anti-inflammatory (NSAID) for the alleviation of pain and inflammation in cattle, sheep, pigs and horses.

<b>Physical Description:</b>	Clear, colourless to yellow liquid.
<b>Schedule:</b>	S4
<b>Active Ingredients:</b>	20 mg/mL MELOXICAM

## WHY CHOOSE USE ILIUM MELOXICAM 20 ANTI-INFLAMMATORY INJECTION?

- ✓ **Injectable formulation** to allow quick administration.
- ✓ **Cost-effective** analgesic, anti-pyretic and anti-inflammatory for use in:
  - Cattle
    - acute respiratory infection and diarrhoea, acute mastitis, and dehorning.
  - Sheep
    - for alleviation of pain and inflammation in sheep and lambs over 14 days of age.
  - Pigs
    - acute non-infectious locomotor disorders, puerperal septicaemia and toxæmia (mastitis, metritis andagalactia syndrome).
  - Horses
    - musculoskeletal disorders and relief of pain associated with colic.
- ✓ **Available in 100 mL and 250 mL pack sizes.**
- ✓ **Plastic packaging** reduces the chance of breakage.
- ✓ **Australian made** which supports local jobs.

# AU PRODUCT INFORMATION SHEET

Information Provided for Veterinary Professionals Only

## ILIMUM MELOXICAM 20

ANTI-INFLAMMATORY INJECTION  
FOR CATTLE, SHEEP, PIGS AND HORSES



### HOW TO USE ILIMUM MELOXICAM 20 ANTI-INFLAMMATORY INJECTION?

#### Dose Rate:

ANIMAL	DOSE
CATTLE	0.5 mg meloxicam/kg body weight i.e. 2.5 mL/100 kg
SHEEP	1.0 mg meloxicam/kg body weight i.e. 1.0 mL / 20 kg
PIGS	0.4 mg meloxicam/kg body weight i.e. 2.0 mL/100 kg Can be repeated once after 24 hours if necessary.
HORSES	0.6 mg meloxicam/kg body weight i.e. 3.0 mL/100 kg Oral meloxicam may be used for continuation of treatment beginning 24 hours after treatment.

#### Administration Route:

Cattle: S.C. or I.V. for single use only.  
Sheep: S.C. for single use only.  
Pigs: I.M.  
Horses: I.V. for single use only.

#### Broach Periods:

ilium Meloxicam 100 mL – 90 days  
ilium Meloxicam 250 mL – 24 hours

#### Precautions:

- Use with caution when using in conjunction with other highly protein bound drugs. The safety of this product in lambs less than 14 days of age has not been established. Use with caution in pregnant sows as safe use has not been fully established. Available information suggests there is no harm when used in the second and third trimester of pregnancy. This product can be used in lactating pigs. Use with caution in pregnant or lactating horses, and in horses under 6 weeks of age as the safety of the product has not been established.
- Concurrent administration with glucocorticosteroids, other non-steroidal anti-inflammatory drugs or with anti-coagulant agents is contraindicated.

#### Contraindications:

- Contraindicated for use in animals suffering from haemorrhagic gastrointestinal disorders, impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of individual hypersensitivity to the product.

**Overdose:** In the case of overdose, symptomatic treatment should be initiated.

#### Withholding Periods:

**Meat:** Cattle: 8 days, Sheep: 11 days, Pigs: 4 days, Horses: 28 days

**Milk:** Cattle: 6 days (12 milkings). Do not feed milk to bobby calves. Do not use in pregnant ewes or lactating ewes within 11 days of lambing where milk may be used or processed for human consumption.

**CATTLE and SHEEP:** Repeat treatments (more than ONCE), higher doses or injection into the muscle could result in residues above the MRLs unless the label withholding period is extended.

**PIGS:** Repeat doses (more than TWICE) or higher doses could result in residues above the MRLs unless the label withholding period is extended.

#### Trade Advice - Export Slaughter Interval (ESI):

**CATTLE:** DO NOT USE less than 21 days before slaughter for export. **SHEEP:** DO NOT USE less than 11 days before slaughter for export. **PIGS:** DO NOT USE less than 4 days before slaughter for export.

**Storage Conditions:** Store below 30 °C.

**Please refer to carton and/or label for full product information.**