

AU PRODUCT INFORMATION SHEET

Information Provided for Veterinary Professionals Only

ilium DERMOTIC EAR & SKIN SUSPENSION



WHAT IS ILIUM DERMOTIC?

It is for the treatment of otitis externa and skin infections caused by fungi, yeasts, Gram-negative and Gram-positive bacteria in dogs and cats. Ilium Dermotic is not active against ear mites.

Physical Description: Liquid suspension

Schedule: S4

Active Ingredients: 23 mg/mL MICONAZOLE NITRATE, 5 mg/mL PREDNISOLONE ACETATE, 0.696 mg/mL POLYMYXIN B SULFATE

WHY CHOOSE ILIUM DERMOTIC?

Otitis Externa is characterised by inflammation of the external ear canal and is most commonly seen in dogs with pendulous ears. Otitis Externa can result from a number of different pathogens.

- ✓ Ilium Dermotic Ear & Skin Suspension contains miconazole, polymyxin B and prednisolone: the synergistic combination addresses the majority of pathogens that cause Otitis Externa as well as treating associated inflammation.
- ✓ **Broad spectrum activity**, antibacterial and anti-fungal
 - Miconazole nitrate and polymyxin B sulfate are antimicrobial
 - Prednisolone is an anti-inflammatory agent.
- ✓ **Proven safety profile.**
- ✓ **Well-tolerated.**
- ✓ **No known interaction** with other medication.
- ✓ **Suspension** provides prolonged contact and distribution within the ear canal.
- ✓ **Formulated for easy administration** & owner compliance.
- ✓ **Australian made.**
- ✓ **Available in 20 mL and 40 mL.**

TR39/63_112A_CAB_112



www.troylab.com.au



(61) 2 8808 3600



admin@troylab.com.au



TROY

Animal
Healthcare

**AUSTRALIAN MADE
ESTABLISHED 1958**

AU PRODUCT INFORMATION SHEET

Information Provided for Veterinary Professionals Only

ilium DERMOTIC

EAR & SKIN SUSPENSION



HOW TO USE ILIUM DERMOTIC?

Shake well before use.

Dose Rate:

Otitis Externa:

- Instil 3 to 5 drops into the ear twice daily.
- Gently massage the ear to ensure even distribution of the product.

Skin infections:

- At the start of treatment, cut the hair surrounding or covering lesions.
- Apply to the affected area and rub in.
- After the clinical signs have completely disappeared, continue treatment without interruption for several days.
- In certain obstinate cases it may be necessary to continue the treatment for 2 to 3 weeks.

Administration Route: Aural and topical.

Contraindications: Do not use where there is a perforated eardrum. Infection or certain procedures might render the tympanic membrane more permeable.

Storage Conditions: Store below 25 °C (air conditioning).

Please refer to carton and/or enclosed leaflet for full product information.

