


NZ PRODUCT INFORMATION SHEET

Information Provided for Veterinary Professionals Only

ILIUM MELOXICAM 20

ANTI-INFLAMMATORY INJECTION
FOR CATTLE, SHEEP, PIGS AND HORSES



WHAT IS ILIUM MELOXICAM 20 ANTI-INFLAMMATORY INJECTION FOR CATTLE, SHEEP, PIGS AND HORSES?

A non-steroidal anti-inflammatory analgesic-antipyretic for use in cattle, sheep, pigs and horses.

Physical Description: Clear, colourless to yellow liquid

Prescription Animal Remedy Class: Restricted Veterinary Medicine

Active Ingredient: 20 mg/mL MELOXICAM

WHY CHOOSE ILIUM MELOXICAM 20 ANTI-INFLAMMATORY INJECTION FOR CATTLE, SHEEP, PIGS AND HORSES?


- ✓ **Injection application** for quick administration.
- ✓ **Cost effective** analgesic, anti-pyretic and anti-inflammatory for use in
 - Cattle:
 - acute respiratory infection and diarrhoea, acute mastitis, and dehorning.
 - Sheep:
 - for alleviation of pain and inflammation in sheep and lambs over 14 days of age.
 - Pigs:
 - acute non-infectious locomotor disorders, puerperal septicaemia and toxæmia (mastitis, metritis and agalactia syndrome).
 - Horses:
 - musculoskeletal disorders and relief of pain associated with colic.
- ✓ **90-day broach period.**
- ✓ **Available in 100 mL plastic container.**
- ✓ **Australian made.**

NZ PRODUCT INFORMATION SHEET

Information Provided for Veterinary Professionals Only

ILIUM MELOXICAM 20

ANTI-INFLAMMATORY INJECTION
FOR CATTLE, SHEEP, PIGS AND HORSES



HOW TO USE ILIUM MELOXICAM 20 ANTI-INFLAMMATORY INJECTION FOR CATTLE, SHEEP, PIGS AND HORSES?

Dose Rate:

ANIMAL	DOSE	ADMINISTRATION ROUTE
 CATTLE	0.5 mg meloxicam/kg body weight i.e. 2.5 mL/100 kg For very young calves weighing less than 50 kg dose at 0.5 mL/20 kg.	S.C. or I.V. for single use only.
 SHEEP	1.0 mg meloxicam/kg body weight i.e. 1.0 mL/20 kg	S.C. for single use only.
 PIGS	0.4 mg meloxicam/kg body weight i.e. 2.0 mL/100 kg Can be repeated once after 24 hours if necessary.	I.M. for single use only.
 HORSES	0.6 mg meloxicam/kg body weight i.e. 3.0 mL/100 kg Oral meloxicam may be used for continuation of treatment beginning 24 hours after treatment.	I.V. for single use only.

Contraindications:

- Contraindicated for use in animals suffering from haemorrhagic gastrointestinal disorders, impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of individual hypersensitivity to the product.
- Concurrent administration with glucocorticosteroids, other non-steroidal anti-inflammatory drugs or with anti-coagulant agents is contraindicated.

Precautions:

- Use with caution when using in conjunction with other highly protein bound drugs. The safety of this product in lambs less than 14 days of age has not been established. Use with caution in pregnant sows as safe use has not been fully established. Available information suggests there is no harm when used in the second and third trimester of pregnancy. This product can be used in lactating pigs. Use with caution in pregnant or lactating horses, and in horses under 6 weeks of age as the safety of the product has not been established.

Overdose: In the case of overdose, symptomatic treatment should be initiated.

Withholding Periods:

Meat: Cattle: 10 days, Sheep: 11 days, Pigs: 3 days, Horses: 28 days

Milk: Cattle: 84 hours, Sheep: Do not use in pregnant ewes or lactating ewes within 11 days of lambing where milk may be used or processed for human consumption.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Storage Conditions: Store below 30 °C (room temperature). Protect from light.

Please refer to carton and/or enclosed leaflet for full product information.