


# NZ PRODUCT INFORMATION SHEET

Information Provided for Veterinary Professionals Only

**ilium**  
**Tolfejec**  
ANTI-INFLAMMATORY INJECTION  
FOR CATTLE AND PIGS



## WHAT IS ILIUM TOLFEJEC ANTI-INFLAMMATORY INJECTION?

Ilium Tolfejec is a non steroidal anti-inflammatory analgesic - antipyretic for use in cattle and pigs.

**Physical Description:** A clear, colourless/pale yellow solution free of visible particulates.

**Prescription Animal Remedy Class:** Restricted Veterinary Medicine

**Active Ingredients:** 40 mg/mL TOLFENAMIC ACID

## WHY CHOOSE ILIUM TOLFEJEC ANTI-INFLAMMATORY INJECTION?


- ✓ **Cost-effective aid** in the control of acute inflammation associated with respiratory disease and as an adjunct in the treatment of acute mastitis in cattle and as an adjunct in the treatment of metritis-mastitis-agalacia in pigs.
- ✓ **Convenient pack size of 100 mL.**
- ✓ **6-month broach period** which allows for plenty of time to use the product and reduces the chance that product will be wasted.
- ✓ **Australian made.**

# NZ PRODUCT INFORMATION SHEET

Information Provided for Veterinary Professionals Only



**ilium**  
**Tolfejec**

ANTI-INFLAMMATORY INJECTION  
FOR CATTLE AND PIGS



## HOW TO USE ILIUM TOLFEJEC ANTI-INFLAMMATORY INJECTION?

### Dose Rate:

 <b>CATTLE</b>	<b>Pneumonia:</b> 2 mg/kg (1 mL per 20 kg b.w.) by intramuscular injection high in the neck (see leaflet for diagram). Treatment may be repeated once only after 48 hours. <b>Mastitis:</b> 4 mg/kg (1 mL per 10 kg b.w.) as a single intravenous injection.
 <b>PIGS</b>	Metritis-mastitis-agalactia: 2 mg/kg (1 mL per 20 kg b.w.) as a single intramuscular injection into the rump.

### Injection route:

I.M. or I.V. dependent on disease.  
Do not exceed 20mL per injection site.

This product may be used during pregnancy and lactation.

### Restraint:

DO NOT inject cattle other than into muscle tissue high on the side of the neck (see product leaflet for diagram). Injection of product into muscles other than as described is likely to result in residues in meat above the MRL.

### Precautions:

Do not exceed the stated dosage and duration of the treatment.

Use aseptic precautions when administering the product.

Do not administer other NSAIDs concurrently or within 24 hours of each other. Tolfenamic acid is highly bound to plasma proteins and may compete with other highly bound drugs.

### Withholding Period:

- MEAT: 14 days.
- MILK: Zero (0) Days.

### Storage:

Store below 30 °C (room temperature). Protect from light.

**Please refer to carton and/or leaflet for full product information.**